

**ANNEX I**  
**SUMMARY OF PRODUCT CHARACTERISTICS**

▼ This medicinal product is subject to additional monitoring. This will allow quick identification of new safety information. Healthcare professionals are asked to report any suspected adverse reactions. See section 4.8 for how to report adverse reactions.

## **1. NAME OF THE MEDICINAL PRODUCT**

Erleada 60 mg film-coated tablets

## **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each film-coated tablet contains 60 mg of apalutamide.

For the full list of excipients, see section 6.1.

## **3. PHARMACEUTICAL FORM**

Film-coated tablet (tablet).

Slightly yellowish to greyish green, oblong-shaped, film-coated tablets (16.7 mm long x 8.7 mm wide), debossed with “AR 60” on one side.

## **4. CLINICAL PARTICULARS**

### **4.1 Therapeutic indications**

Erleada is indicated:

- in adult men for the treatment of non-metastatic castration-resistant prostate cancer (nmCRPC) who are at high risk of developing metastatic disease (see section 5.1).
- in adult men for the treatment of metastatic hormone-sensitive prostate cancer (mHSPC) in combination with androgen deprivation therapy (ADT) (see section 5.1).

### **4.2 Posology and method of administration**

Treatment with apalutamide should be initiated and supervised by specialist physicians experienced in the medical treatment of prostate cancer.

#### Posology

The recommended dose is 240 mg (four 60 mg tablets) as an oral single daily dose.

Medical castration with gonadotropin releasing hormone analogue (GnRHa) should be continued during treatment in patients not surgically castrated.

If a dose is missed, it should be taken as soon as possible on the same day with a return to the normal schedule the following day. Extra tablets should not be taken to make up the missed dose.

If a  $\geq$  Grade 3 toxicity or an intolerable adverse reaction is experienced by the patient, dosing should be held rather than permanently discontinuing treatment until symptoms improve to  $\leq$  Grade 1 or original grade, then should be resumed at the same dose or a reduced dose (180 mg or 120 mg), if warranted. For the most common adverse reactions, see section 4.8.

## *Special populations*

### Elderly

No dose adjustment is necessary for elderly patients (see sections 5.1 and 5.2).

### Renal impairment

No dose adjustment is necessary for patients with mild to moderate renal impairment.

Caution is required in patients with severe renal impairment as apalutamide has not been studied in this patient population (see section 5.2). If treatment is started, patients should be monitored for the adverse reactions listed in section 4.8 and dose reduce as per section 4.2 Posology and method of administration.

### Hepatic impairment

No dose adjustment is necessary for patients with baseline mild or moderate hepatic impairment (Child-Pugh Class A and B, respectively).

Erleada is not recommended in patients with severe hepatic impairment as there are no data in this patient population and apalutamide is primarily hepatically eliminated (see section 5.2).

### Paediatric population

There is no relevant use of apalutamide in the paediatric population.

### Method of administration

Oral use.

The tablets should be swallowed whole and can be taken with or without food.

## **4.3 Contraindications**

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

Women who are or may become pregnant (see section 4.6).

## **4.4 Special warnings and precautions for use**

### Seizure

Erleada is not recommended in patients with a history of seizures or other predisposing factors including, but not limited to, underlying brain injury, recent stroke (within one year), primary brain tumours or brain metastases. If a seizure develops during treatment with Erleada, treatment should be discontinued permanently. The risk of seizure may be increased in patients receiving concomitant medicinal products that lower the seizure threshold.

In two randomised studies (SPARTAN and TITAN), seizure occurred in 0.4% of patients receiving apalutamide and in 0.2% of patients treated with placebo. These studies excluded patients with a history of seizure or predisposing factors for seizure.

There is no clinical experience in re-administering Erleada to patients who experienced a seizure.

### Falls and fractures

Falls and fractures occurred in patients receiving apalutamide (see section 4.8). Patients should be evaluated for fracture and fall risk before starting Erleada and should continue to be monitored and managed according to established treatment guidelines and use of bone-targeted agents should be considered.

### Ischaemic heart disease

Ischaemic heart disease, including events leading to death, occurred in patients treated with apalutamide (see section 4.8). The majority of patients had cardiac risk factors. Patients should be monitored for signs and symptoms of ischaemic heart disease and management of cardiovascular risk factors, such as hypertension, diabetes, or dyslipidaemia should be optimised as per standard of care.

### Concomitant use with other medicinal products

Apalutamide is a potent enzyme inducer and may lead to loss of efficacy of many commonly used medicinal products (see section 4.5). A review of concomitant medicinal products should therefore be conducted when apalutamide treatment is initiated. Concomitant use of apalutamide with medicinal products that are sensitive substrates of many metabolising enzymes or transporters (see section 4.5) should generally be avoided if their therapeutic effect is of large importance to the patient, and if dose adjustments cannot easily be performed based on monitoring of efficacy or plasma concentrations.

Co-administration of apalutamide with warfarin and coumarin-like anticoagulants should be avoided. If Erleada is co-administered with an anticoagulant metabolised by CYP2C9 (such as warfarin or acenocoumarol), additional International Normalised Ratio (INR) monitoring should be conducted (see section 4.5).

### Recent cardiovascular disease

Patients with clinically significant cardiovascular disease in the past 6 months including severe/unstable angina, myocardial infarction, symptomatic congestive heart failure, arterial or venous thromboembolic events (e.g., pulmonary embolism, cerebrovascular accident including transient ischaemic attacks), or clinically significant ventricular arrhythmias were excluded from the clinical studies. Therefore, the safety of apalutamide in these patients has not been established. If Erleada is prescribed, patients with clinically significant cardiovascular disease should be monitored for risk factors such as hypercholesterolaemia, hypertriglyceridaemia, or other cardio-metabolic disorders (see section 4.8). Patients should be treated, if appropriate, after initiating Erleada for these conditions according to established treatment guidelines.

### Androgen deprivation therapy may prolong the QT interval

In patients with a history of or risk factors for QT prolongation and in patients receiving concomitant medicinal products that might prolong the QT interval (see section 4.5), physicians should assess the benefit-risk ratio including the potential for Torsade de pointes prior to initiating Erleada.

## **4.5 Interaction with other medicinal products and other forms of interaction**

The elimination of apalutamide and formation of its active metabolite, N-desmethyl apalutamide, is mediated by both CYP2C8 and CYP3A4 to a similar extent at steady-state. No clinically meaningful changes in their overall exposure is expected as a result of drug interaction with inhibitors or inducers of CYP2C8 or CYP3A4. Apalutamide is an inducer of enzymes and transporters and may lead to an increase in elimination of many commonly used medicinal products.

### Potential for other medicinal products to affect apalutamide exposures

#### *Medicinal products that inhibit CYP2C8*

CYP2C8 plays a role in the elimination of apalutamide and in the formation of its active metabolite. In a drug-drug interaction study, the  $C_{max}$  of apalutamide decreased by 21% while AUC increased by 68% following co-administration of apalutamide 240 mg single dose with gemfibrozil (strong CYP2C8 inhibitor). For the active moieties (sum of apalutamide plus the potency adjusted active metabolite),  $C_{max}$  decreased by 21% while AUC increased by 45%. No initial dose adjustment is necessary when Erleada is co-administered with a strong inhibitor of CYP2C8 (e.g., gemfibrozil, clopidogrel) however, a reduction of the Erleada dose based on tolerability should be considered (see section 4.2). Mild or moderate inhibitors of CYP2C8 are not expected to affect the exposure of apalutamide.

#### *Medicinal products that inhibit CYP3A4*

CYP3A4 plays a role in the elimination of apalutamide and in the formation of its active metabolite. In a drug-drug interaction study, the  $C_{max}$  of apalutamide decreased by 22% while AUC was similar following co-administration of Erleada as a 240 mg single dose with itraconazole (strong CYP3A4 inhibitor). For the active moieties (sum of apalutamide plus the potency adjusted active metabolite),  $C_{max}$  decreased by 22% while AUC was again similar. No initial dose adjustment is necessary when Erleada is co-administered with a strong inhibitor of CYP3A4 (e.g., ketoconazole, ritonavir, clarithromycin) however, a reduction of the Erleada dose based on tolerability should be considered (see section 4.2). Mild or moderate inhibitors of CYP3A4 are not expected to affect the exposure of apalutamide.

#### *Medicinal products that induce CYP3A4 or CYP2C8*

The effects of CYP3A4 or CYP2C8 inducers on the pharmacokinetics of apalutamide have not been evaluated *in vivo*. Based on the drug-drug interaction study results with strong CYP3A4 inhibitor or strong CYP2C8 inhibitor, CYP3A4 or CYP2C8 inducers are not expected to have clinically relevant effects on the pharmacokinetics of apalutamide and the active moieties therefore no dose adjustment is necessary when Erleada is co-administered with inducers of CYP3A4 or CYP2C8.

#### Potential for apalutamide to affect exposures to other medicinal products

Apalutamide is a potent enzyme inducer and increases the synthesis of many enzymes and transporters; therefore, interaction with many common medicinal products that are substrates of enzymes or transporters is expected. The reduction in plasma concentrations can be substantial, and lead to lost or reduced clinical effect. There is also a risk of increased formation of active metabolites.

#### *Drug metabolising enzymes*

*In vitro* studies showed that apalutamide and N-desmethyl apalutamide are moderate to strong CYP3A4 and CYP2B6 inducers, are moderate inhibitors of CYP2B6 and CYP2C8, and weak inhibitors of CYP2C9, CYP2C19, and CYP3A4. Apalutamide and N-desmethyl apalutamide do not affect CYP1A2 and CYP2D6 at therapeutically relevant concentrations. The effect of apalutamide on CYP2B6 substrates has not been evaluated *in vivo* and the net effect is presently unknown. When substrates of CYP2B6 (e.g., efavirenz) are administered with Erleada, monitoring for an adverse reaction and evaluation for loss of efficacy of the substrate should be performed and dose adjustment of the substrate may be required to maintain optimal plasma concentrations.

In humans, apalutamide is a strong inducer of CYP3A4 and CYP2C19, and a weak inducer of CYP2C9. In a drug-drug interaction study using a cocktail approach, co-administration of apalutamide with single oral doses of sensitive CYP substrates resulted in a 92% decrease in the AUC of midazolam (CYP3A4 substrate), 85% decrease in the AUC of omeprazole (CYP2C19 substrate), and 46% decrease in the AUC of S-warfarin (CYP2C9 substrate). Apalutamide did not cause clinically meaningful changes in exposure to the CYP2C8 substrate. Concomitant use of Erleada with medicinal products that are primarily metabolised by CYP3A4 (e.g., darunavir, felodipine, midazolam, simvastatin), CYP2C19 (e.g., diazepam, omeprazole), or CYP2C9 (e.g., warfarin, phenytoin) can result in lower exposure to these medicinal products. Substitution for these medicinal products is recommended when possible or evaluation for loss of efficacy should be performed if the medicinal product is continued. If given with warfarin, INR should be monitored during Erleada treatment.

Induction of CYP3A4 by apalutamide suggests that UDP-glucuronosyl transferase (UGT) may also be induced via activation of the nuclear pregnane X receptor (PXR). Concomitant administration of Erleada with medicinal products that are substrates of UGT (e.g., levothyroxine, valproic acid) can result in lower exposure to these medicinal products. When substrates of UGT are co-administered with Erleada, evaluation for loss of efficacy of the substrate should be performed and dose adjustment of the substrate may be required to maintain optimal plasma concentrations.

#### *Drug transporters*

Apalutamide was shown to be a weak inducer of P-glycoprotein (P-gp), breast cancer resistance protein (BCRP), and organic anion transporting polypeptide 1B1 (OATP1B1) clinically. A drug-drug

interaction study using a cocktail approach showed that co-administration of apalutamide with single oral doses of sensitive transporter substrates resulted in a 30% decrease in the AUC of fexofenadine (P-gp substrate) and 41% decrease in the AUC of rosuvastatin (BCRP/OATP1B1 substrate) but had no impact on  $C_{max}$ . Concomitant use of Erleada with medicinal products that are substrates of P-gp (e.g., colchicine, dabigatran etexilate, digoxin), BCRP or OATP1B1 (e.g., lapatinib, methotrexate, rosuvastatin, repaglinide) can result in lower exposure of these medicinal products. When substrates of P-gp, BCRP or OATP1B1 are co-administered with Erleada, evaluation for loss of efficacy of the substrate should be performed and dose adjustment of the substrate may be required to maintain optimal plasma concentrations.

Based on *in vitro* data, inhibition of organic cation transporter 2 (OCT2), organic anion transporter 3 (OAT3) and multidrug and toxin extrusions (MATEs) by apalutamide and its N-desmethyl metabolite cannot be excluded. No *in vitro* inhibition of organic anion transporter 1 (OAT1) was observed.

#### GnRH Analog

In mHSPC subjects receiving leuprolide acetate (a GnRH analog), co-administration with apalutamide had no apparent effect on the steady-state exposure of leuprolide.

#### Medicinal products which prolong the QT interval

Since androgen deprivation treatment may prolong the QT interval, the concomitant use of Erleada with medicinal products known to prolong the QT interval or medicinal products able to induce Torsade de pointes such as class IA (e.g., quinidine, disopyramide) or class III (e.g., amiodarone, sotalol, dofetilide, ibutilide) antiarrhythmic medicinal products, methadone, moxifloxacin, antipsychotics (e.g. haloperidol), etc. should be carefully evaluated (see section 4.4).

#### Paediatric population

Interaction studies have only been performed in adults.

### **4.6 Fertility, pregnancy and lactation**

#### Contraception in males and females

It is not known whether apalutamide or its metabolites are present in semen. Erleada may be harmful to a developing foetus. For patients having sex with female partners of reproductive potential, a condom should be used along with another highly effective contraceptive method during treatment and for 3 months after the last dose of Erleada.

#### Pregnancy

Erleada is contraindicated in women who are or may become pregnant (see section 4.3). Based on its mechanism of action, Erleada may cause foetal harm when administered during pregnancy. There are no data available from the use of Erleada in pregnant women. Animal reproductive studies have not been conducted with Erleada.

#### Breast-feeding

It is unknown whether apalutamide/metabolites are excreted in human milk. A risk to the suckling child cannot be excluded. Erleada should not be used during breast-feeding.

#### Fertility

Based on animal studies, Erleada may decrease fertility in males of reproductive potential (see section 5.3).

### **4.7 Effects on ability to drive and use machines**

Erleada has no or negligible influence on the ability to drive and use machines. However, seizures have been reported in patients taking Erleada. Patients should be advised of this risk in regards to driving or operating machines.

## 4.8 Undesirable effects

### Summary of the safety profile

The most common adverse reactions are fatigue (26%), skin rash (26% of any grade and 6% Grade 3 or 4), hypertension (22%), hot flush (18%), arthralgia (17%), diarrhoea (16%), fall (13%), and weight decreased (13%). Other important adverse reactions include fractures (11%) and hypothyroidism (8%).

### Tabulated list of adverse reactions

Adverse reactions observed during clinical studies are listed below by frequency category. Frequency categories are defined as follows: very common ( $\geq 1/10$ ); common ( $\geq 1/100$  to  $< 1/10$ ); uncommon ( $\geq 1/1,000$  to  $< 1/100$ ); rare ( $\geq 1/10,000$  to  $< 1/1,000$ ); very rare ( $< 1/10,000$ ) and not known (frequency cannot be estimated from the available data).

Within each frequency grouping, undesirable effects are presented in order of decreasing seriousness.

**Table 1: Adverse reactions identified in clinical studies**

System organ class	Adverse reaction and frequency
<b>Endocrine disorders</b>	common: hypothyroidism <sup>a</sup>
<b>Metabolism and nutrition disorders</b>	very common: decreased appetite
	common: hypercholesterolaemia, hypertriglyceridaemia
<b>Nervous system disorders</b>	common: dysgeusia
	uncommon: seizure <sup>b</sup> (see section 4.4)
<b>Cardiac disorders</b>	common: ischaemic heart disease <sup>c</sup>
	not known: QT prolongation (see sections 4.4 and 4.5)
<b>Vascular disorders</b>	very common: hot flush, hypertension
<b>Gastrointestinal disorders</b>	very common: diarrhoea
<b>Skin and subcutaneous tissue disorders</b>	very common: skin rash <sup>d</sup>
	common: pruritus
	not known: toxic epidermal necrolysis <sup>e</sup>
<b>Musculoskeletal and connective tissue disorders</b>	very common: fracture <sup>f</sup> , arthralgia
	common: muscle spasm
<b>General disorders and administration site conditions</b>	very common: fatigue
<b>Investigations</b>	very common: weight decreased
<b>Injury, poisoning and procedural complications</b>	very common: fall

<sup>a</sup> Includes hypothyroidism, blood thyroid stimulating hormone increased, thyroxine decreased, autoimmune thyroiditis, thyroxine free decreased, tri-iodothyronine decreased

<sup>b</sup> Includes tongue biting

<sup>c</sup> Includes angina pectoris, angina unstable, myocardial infarction, acute myocardial infarction, coronary artery occlusion, coronary artery stenosis, acute coronary syndrome, arteriosclerosis coronary artery, cardiac stress test abnormal, troponin increased, myocardial ischaemia

<sup>d</sup> See "Skin rash" under "Description of selected adverse reactions"

<sup>e</sup> Post-marketing adverse reaction

<sup>f</sup> Includes rib fracture, lumbar vertebral fracture, spinal compression fracture, spinal fracture, foot fracture, hip fracture, humerus fracture, thoracic vertebral fracture, upper limb fracture, fractured sacrum, hand fracture, pubis fracture, acetabulum fracture, ankle fracture, compression fracture, costal cartilage fracture, facial bones fracture, lower limb fracture, osteoporotic fracture, wrist fracture, avulsion fracture, fibula fracture, fractured coccyx, pelvic fracture, radius fracture, sternal fracture, stress fracture, traumatic fracture, cervical vertebral fracture, femoral neck fracture, tibia fracture. See below.

### Description of selected adverse reactions

#### *Skin rash*

Skin rash associated with apalutamide was most commonly described as macular or maculo-papular. Skin rash included rash, rash maculo-papular, rash generalised, urticaria, rash pruritic, rash macular, conjunctivitis, erythema multiforme, rash papular, skin exfoliation, genital rash, rash erythematous,

stomatitis, drug eruption, mouth ulceration, rash pustular, blister, papule, pemphigoid, skin erosion, dermatitis, and rash vesicular. Adverse reactions of skin rash were reported for 26% of patients treated with apalutamide. Grade 3 skin rashes (defined as covering > 30% body surface area [BSA]) were reported with apalutamide treatment in 6% of patients.

The median days to onset of skin rash was 83 days. Seventy-eight percent of patients had resolution of rash with a median of 78 days to resolution. Medicinal products utilised included topical corticosteroids, oral anti-histamines, and 19% of patients received systemic corticosteroids. Among patients with skin rash, dose interruption occurred in 28% and dose reduction occurred in 14% (see section 4.2). Skin rash recurred in 59% of patients who had dose interruption. Skin rash led to apalutamide treatment discontinuation in 7% of patients who experienced skin rash.

#### *Falls and fractures*

In Study ARN-509-003, fracture was reported for 11.7% of patients treated with apalutamide and 6.5% of patients treated with placebo. Half of the patients experienced a fall within 7 days before the fracture event in both treatment groups. Falls were reported for 15.6% of patients treated with apalutamide *versus* 9.0% of patients treated with placebo (see section 4.4).

#### *Ischaemic heart disease*

In a randomised study (SPARTAN) of patients with nmCRPC, ischaemic heart disease occurred in 4% of patients treated with apalutamide and 3% of patients treated with placebo. In a randomised study (TITAN) in patients with mHSPC, ischaemic heart disease occurred in 4% of patients treated with apalutamide and 2% of patients treated with placebo. Across the SPARTAN and TITAN studies, 6 patients (0.5%) treated with apalutamide and 2 patients (0.2%) treated with placebo died from an ischaemic heart disease (see section 4.4).

#### *Hypothyroidism*

Hypothyroidism was reported for 8% of patients treated with apalutamide and 2% of patients treated with placebo based on assessments of thyroid-stimulating hormone (TSH) every 4 months. There were no grade 3 or 4 adverse events. Hypothyroidism occurred in 30% of patients already receiving thyroid replacement therapy in the apalutamide arm and in 3% of patients in the placebo arm. In patients not receiving thyroid replacement therapy, hypothyroidism occurred in 7% of patients treated with apalutamide and in 2% of patients treated with placebo. Thyroid replacement therapy, when clinically indicated, should be initiated or dose-adjusted (see section 4.5).

#### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via [the national reporting system listed in Appendix V](#).

## **4.9 Overdose**

There is no known specific antidote for apalutamide overdose. In the event of an overdose, Erleada should be stopped and general supportive measures should be undertaken until clinical toxicity has been diminished or resolved. Adverse reactions in the event of an overdose has not yet been observed, it is expected that such reactions would resemble the adverse reactions listed in section 4.8.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Endocrine therapy, anti-androgens, ATC code: L02BB05



### Mechanism of action

Apalutamide is an orally administered, selective Androgen Receptor (AR) inhibitor that binds directly to the ligand-binding domain of the AR. Apalutamide prevents AR nuclear translocation, inhibits DNA binding, impedes AR-mediated transcription, and lacks androgen receptor agonist activity. Apalutamide treatment decreases tumor cell proliferation and increases apoptosis leading to potent antitumor activity. A major metabolite, N-desmethyl apalutamide, exhibited one-third the *in vitro* activity of apalutamide.

### Cardiac electrophysiology

The effect of apalutamide 240 mg once daily on the QTc interval was assessed in an open-label, uncontrolled, multi-center, single-arm dedicated QT study in 45 patients with CRPC. At steady-state, the maximum mean QTcF change from baseline was 12.4 ms (2-sided 90% upper CI: 16.0 ms). An exposure-QT analysis suggested a concentration-dependent increase in QTcF for apalutamide and its active metabolite.

### Clinical efficacy and safety

The efficacy and safety of apalutamide has been established in two Phase 3 randomised, placebo-controlled studies, Study ARN-509-003 (nmCRPC) and 56021927PCR3002 (mHSPC).

### TITAN: Metastatic Hormone-sensitive Prostate Cancer (mHSPC)

TITAN was a randomised, double-blind, placebo-controlled, multinational, multicenter clinical trial in which 1052 patients with mHSPC were randomised (1:1) to receive either apalutamide orally at a dose of 240 mg once daily (N = 525) or placebo once daily (N = 527). All patients were required to have at least one bone metastasis on Technetium <sup>99m</sup> bone scan. Patients were excluded if the site of metastases was limited to either the lymph nodes or viscera (e.g., liver or lung). All patients in the TITAN trial received concomitant GnRH analog or had prior bilateral orchiectomy. Around 11% of patients received prior treatment with docetaxel (maximum of 6 cycles, last dose ≤2 months prior to randomisation and maintained response prior to randomisation). The exclusion criteria included known brain metastases; prior treatment with other next generation anti-androgens (eg, enzalutamide), CYP17 inhibitors (eg, abiraterone acetate), immunotherapy (eg, sipuleucel-T), radiopharmaceutical agents or other treatments for prostate cancer; or history of seizure or condition that may predispose to seizure. Patients were stratified by Gleason score at diagnosis, prior docetaxel use, and region of the world. Patients with both high- and low-volume mHSPC were eligible for the study. High-volume disease was defined as either visceral metastases and at least 1 bone lesion or at least 4 bone lesions, with at least 1 bone lesion outside of the vertebral column or pelvis. Low-volume disease was defined as the presence of bone lesion(s) not meeting the definition of high-volume.

The following patient demographics and baseline disease characteristics were balanced between the treatment arms. The median age was 68 years (range 43-94) and 23% of patients were 75 years of age or older. The racial distribution was 68% Caucasian, 22% Asian, and 2% Black. Sixty-three percent (63%) of patients had high-volume disease and 37% had low-volume disease. Sixteen percent (16%) of patients had prior surgery, radiotherapy of the prostate or both. A majority of patients had a Gleason score of 7 or higher (92%). Sixty-eight percent (68%) of patients received prior treatment with a first-generation anti-androgen in the non-metastatic setting. Although criteria for castration resistance were not determined at baseline, 94% of patients demonstrated a decrease in prostate specific antigen (PSA) from initiation of androgen deprivation therapy (ADT) to first dose of apalutamide or placebo. All patients except one in the placebo group, had an Eastern Cooperative Oncology Group Performance Status (ECOG PS) score of 0 or 1 at study entry. Among the patients who discontinued study treatment (N = 271 for placebo and N = 170 for Erleada), the most common reason for discontinuation in both arms was disease progression. A greater proportion (73%) of patients treated with placebo received subsequent anti-cancer therapy compared to patients treated with Erleada (54%).

The major efficacy outcome measures of the study were overall survival (OS) and radiographic progression-free survival (rPFS). Efficacy results of TITAN are summarised in Table 2 and Figures 1 and 2.

**Table 2: Summary of Efficacy Results – Intent-to-treat mHSPC Population (TITAN)**

Endpoint	Erleada N=525	Placebo N=527
<b>Overall Survival</b>		
Deaths (%)	83 (16%)	117 (22%)
Median, months (95% CI)	NE (NE, NE)	NE (NE, NE)
Hazard ratio (95% CI) <sup>a</sup>	0.671 (0.507, 0.890)	
p-value <sup>b</sup>	0.0053	
<b>Radiographic Progression-free Survival</b>		
Disease progression or death (%)	134 (26%)	231 (44%)
Median, months (95% CI)	NE (NE, NE)	22.08 (18.46, 32.92)
Hazard ratio (95% CI) <sup>a</sup>	0.484 (0.391, 0.600)	
p-value <sup>b</sup>	<.0001	


<sup>a</sup> Hazard ratio is from stratified proportional hazards model. Hazard ratio <1 favors active treatment.

<sup>b</sup> p-value is from the log-rank test stratified by Gleason score at diagnosis ( $\leq 7$  vs.  $>7$ ), Region (NA/EU vs. Other Countries) and Prior docetaxel use (Yes vs. No).


NE=Not Estimable

A statistically significant improvement in OS and rPFS was demonstrated in patients randomised to receive Erleada compared with patients randomised to receive placebo. Consistent improvement in rPFS was observed across patient subgroups including high- or low-volume disease, prior docetaxel use (yes or no), age ( $< 65$ ,  $\geq 65$ , or  $\geq 75$  years old), baseline PSA above median (yes or no), and number of bone lesions ( $\leq 10$  or  $>10$ ).

**Figure 1: Kaplan-Meier Plot of Overall Survival (OS); Intent-to-treat mHSPC Population (TITAN)**



**Figure 2: Kaplan-Meier Plot of Radiographic Progression-Free Survival (rPFS); Intent-to-treat mHSPC Population (TITAN)**



Treatment with Erleada statistically significantly delayed the initiation of cytotoxic chemotherapy (HR = 0.391, CI = 0.274, 0.558; p < 0.0001), resulting in a 61% reduction of risk for subjects in the treatment arm compared to the placebo arm.

***SPARTAN: Non-Metastatic Castration Resistant Prostate Cancer (nmCRPC)***


A total of 1207 subjects with NM-CRPC were randomised 2:1 to receive either apalutamide orally at a dose of 240 mg once daily in combination with androgen deprivation therapy (ADT) (medical castration or prior surgical castration) or placebo with ADT in a multicenter, double-blind, clinical study (Study ARN-509-003). Subjects enrolled had a Prostate Specific Antigen (PSA) Doubling Time (PSADT) ≤ 10 months, considered to be at high risk of imminent metastatic disease and prostate cancer-specific death. All subjects who were not surgically castrated received ADT continuously throughout the study. PSA results were blinded and were not used for treatment discontinuation. Subjects randomised to either arm were to continue treatment until disease progression defined by blinded central imaging review (BICR), initiation of new treatment, unacceptable toxicity or withdrawal.

The following patient demographics and baseline disease characteristics were balanced between the treatment arms. The median age was 74 years (range 48-97) and 26% of subjects were 80 years of age or older. The racial distribution was 66% Caucasian, 5.6% Black, 12% Asian, and 0.2% Other. Seventy-seven percent (77%) of subjects in both treatment arms had prior surgery or radiotherapy of the prostate. A majority of subjects had a Gleason score of 7 or higher (81%). Fifteen percent (15%) of subjects had < 2 cm pelvic lymph nodes at study entry. Seventy-three percent (73%) of subjects received prior treatment with a first generation anti-androgen; 69% of subjects received bicalutamide and 10% of subjects received flutamide. All subjects enrolled were confirmed to be non-metastatic by

blinded central imaging review and had an Eastern Cooperative Oncology Group Performance Status (ECOG PS) performance status score of 0 or 1 at study entry.

Metastasis-free survival (MFS) was the primary endpoint, defined as the time from randomisation to the time of first evidence of BICR-confirmed bone or soft tissue distant metastasis or death due to any cause, whichever occurred first. Treatment with Erleada significantly improved MFS. Erleada decreased the relative risk of distant metastasis or death by 70% compared to placebo (HR = 0.30; 95% CI: 0.24, 0.36;  $p < 0.0001$ ). The median MFS for Erleada was 41 months and was 16 months for placebo (see Figure 3). Consistent improvement in MFS with Erleada was observed for all pre-specified subgroups, including age, race, region of the world, nodal status, prior number of hormonal therapies, baseline PSA, PSA doubling time, baseline ECOG status and use of bone-sparing agents.

**Figure 3: Kaplan-Meier metastasis-free survival (MFS) curve in Study ARN-509-003**



Subjects treated with Erleada and ADT showed significant improvement over those treated with ADT alone for the following secondary endpoints of time to metastasis (HR = 0.28; 95% CI: 0.23, 0.34;  $p < 0.0001$ ), progression-free survival (PFS) (HR = 0.30; 95% CI: 0.25, 0.36;  $p < 0.0001$ ); time to symptomatic progression (HR = 0.45; 95% CI: 0.32, 0.63;  $p < 0.0001$ ); and a trend for overall survival (OS) (HR = 0.70; 95% CI: 0.47 1.04;  $p = 0.0742$ ).

Time to symptomatic progression was defined as time from randomization to development of a skeletal related event, pain/symptoms requiring initiation of a new systemic anti-cancer therapy, or loco-regional tumor progression requiring radiation/surgery. While the overall number of events was small, the difference between the two arms was sufficiently large to reach statistical significance. In the apalutamide arm 64 (7.9%) subjects developed symptomatic progression compared with 63 (16%) in the placebo arm, with a hazard ratio of 0.447 (95% CI: 0.315, 0.634), which met the pre-specified O'Brien Fleming type efficacy boundary for significance,  $p < 0.00008$ . The median time to symptomatic progression was not reached in either treatment group.

In the apalutamide arm, 62 (7.7%) subjects died compared to 42 (10.5%) subjects in the placebo arm. The median survival for the apalutamide arm was not reached compared to 39.03 months with a 95%

CI of (39.03, NE) for the placebo arm. Statistical significance was not reached in overall survival at the pre-specified interim analysis.

Post-progression survival (PFS-2, defined as the time to disease progression after first subsequent therapy or death) was longer for subjects treated with Erleada compared to those treated with placebo (HR = 0.489; 95% CI: 0.361, 0.662;  $p < 0.0001$ ).

There were no statistically significant differences observed in the change from baseline analysis of the Functional Assessment of Cancer Therapy-Prostate (FACT-P) for the total score or any of the subscales between subjects on Erleada added to ADT *versus* placebo with ADT.

#### Paediatric population

The European Medicines Agency has waived the obligation to submit the results of studies with Erleada in all subsets of the paediatric population in advanced prostate cancer. See section 4.2 for information on paediatric use.

## **5.2 Pharmacokinetic properties**

Following repeat once-daily dosing, apalutamide exposure ( $C_{max}$  and area under the concentration curve [AUC]) increased in a dose-proportional manner across the dose range of 30 to 480 mg. Following administration of 240 mg once daily, apalutamide steady state was achieved after 4 weeks and the mean accumulation ratio was approximately 5-fold relative to a single dose. At steady-state, mean (CV%)  $C_{max}$  and AUC values for apalutamide were 6  $\mu\text{g/mL}$  (28%) and 100  $\mu\text{g.h/mL}$  (32%), respectively. Daily fluctuations in apalutamide plasma concentrations were low, with mean peak-to-trough ratio of 1.63. An increase in apparent clearance ( $CL/F$ ) was observed with repeat dosing, likely due to induction of apalutamide's own metabolism.

At steady-state, the mean (CV%)  $C_{max}$  and AUC values for the major active metabolite, N-desmethyl apalutamide, were 5.9  $\mu\text{g/mL}$  (18%) and 124  $\mu\text{g.h/mL}$  (19%), respectively. N-desmethyl apalutamide is characterised by a flat concentration-time profile at steady-state with a mean peak-to-trough ratio of 1.27. Mean (CV%) AUC metabolite/parent drug ratio for N-desmethyl apalutamide following repeat-dose administration was about 1.3 (21%). Based on systemic exposure, relative potency, and pharmacokinetic properties, N-desmethyl apalutamide likely contributed to the clinical activity of apalutamide.

#### Absorption

After oral administration, median time to achieve peak plasma concentration ( $t_{max}$ ) was 2 hours (range: 1 to 5 hours). Mean absolute oral bioavailability is approximately 100%, indicating that apalutamide is completely absorbed after oral administration.

Administration of apalutamide to healthy subjects under fasting conditions and with a high-fat meal resulted in no clinically relevant changes in  $C_{max}$  and AUC. Median time to reach  $t_{max}$  was delayed about 2 hours with food (see section 4.2).

Apalutamide is not ionizable under relevant physiological pH condition, therefore acid lowering agents (e.g., proton pump inhibitor,  $H_2$ -receptor antagonist, antacid) are not expected to affect the solubility and bioavailability of apalutamide.

*In vitro*, apalutamide and its N-desmethyl metabolite are substrates for P-gp. Because apalutamide is completely absorbed after oral administration, P-gp does not limit the absorption of apalutamide and therefore, inhibition or induction of P-gp is not expected to affect the bioavailability of apalutamide.

#### Distribution

The mean apparent volume of distribution at steady-state of apalutamide is about 276 L. The volume of distribution of apalutamide is greater than the volume of total body water, indicative of extensive extravascular distribution.

Apalutamide and N-desmethyl apalutamide are 96% and 95% bound to plasma proteins, respectively, and mainly bind to serum albumin with no concentration dependency.

#### Biotransformation

Following single oral administration of <sup>14</sup>C-labeled apalutamide 240 mg, apalutamide, the active metabolite, N-desmethyl apalutamide, and an inactive carboxylic acid metabolite accounted for the majority of the <sup>14</sup>C-radioactivity in plasma, representing 45%, 44%, and 3%, respectively, of the total <sup>14</sup>C-AUC.

Metabolism is the main route of elimination of apalutamide. It is metabolised primarily by CYP2C8 and CYP3A4 to form N-desmethyl apalutamide. Apalutamide and N-desmethyl apalutamide are further metabolised to form the inactive carboxylic acid metabolite by carboxylesterase. The contribution of CYP2C8 and CYP3A4 in the metabolism of apalutamide is estimated to be 58% and 13% following single dose but the level of contribution is expected to change at steady-state due to induction of CYP3A4 by apalutamide after repeat dose.

#### Elimination

Apalutamide, mainly in the form of metabolites, is eliminated primarily via urine. Following a single oral administration of radiolabeled apalutamide, 89% of the radioactivity was recovered up to 70 days post-dose: 65% was recovered in urine (1.2% of dose as unchanged apalutamide and 2.7% as N-desmethyl apalutamide) and 24% was recovered in feces (1.5% of dose as unchanged apalutamide and 2% as N-desmethyl apalutamide).

The apparent oral clearance (CL/F) of apalutamide is 1.3 L/h after single dosing and increases to 2.0 L/h at steady-state after once-daily dosing. The mean effective half-life for apalutamide in patients is about 3 days at steady-state.

*In vitro* data indicate that apalutamide and its N-desmethyl metabolite are not substrates for BCRP, OATP1B1 or OATP1B3.

#### Special populations

The effects of renal impairment, hepatic impairment, age, race, and other extrinsic factors on the pharmacokinetics of apalutamide are summarised below.

#### Renal impairment

A dedicated renal impairment study for apalutamide has not been conducted. Based on the population pharmacokinetic analysis using data from clinical studies in subjects with castration-resistant prostate cancer (CRPC) and healthy subjects, no significant difference in systemic apalutamide exposure was observed in subjects with pre-existing mild to moderate renal impairment (estimated glomerular filtration rate [eGFR] between 30 to 89 mL/min/1.73 m<sup>2</sup>; N=585) compared to subjects with baseline normal renal function (eGFR ≥ 90 mL/min/1.73 m<sup>2</sup>; N=372). The potential effect of severe renal impairment or end stage renal disease (eGFR ≤ 29 mL/min/1.73 m<sup>2</sup>) have not been established due to insufficient data.

#### Hepatic impairment

A dedicated hepatic impairment study compared the systemic exposure of apalutamide and N-desmethyl apalutamide in subjects with baseline mild hepatic impairment (N=8, Child-Pugh Class A, mean score = 5.3) or moderate hepatic impairment (N=8, Child-Pugh Class B, mean score = 7.6) *versus* healthy controls with normal hepatic function (N=8). Following a single oral 240 mg dose of apalutamide, the geometric mean ratio (GMR) for AUC and C<sub>max</sub> for apalutamide in subjects with mild impairment was 95% and 102%, respectively, and the GMR for AUC and C<sub>max</sub> of apalutamide in subjects with moderate impairment was 113% and 104%, respectively, compared to healthy control subjects. Clinical and pharmacokinetic data for apalutamide are not available for patients with severe hepatic impairment (Child-Pugh Class C).

### Ethnicity and race

Based on population pharmacokinetic analysis, there were no clinically relevant differences in apalutamide pharmacokinetics between White (Caucasian or Hispanic or Latino; N=761), Black (of African heritage or African American; N=71), Asian (non-Japanese; N=58) and Japanese (N=58).

### Age

Population pharmacokinetic analyses showed that age (range: 18 to 94 years) does not have a clinically meaningful influence on the pharmacokinetics of apalutamide.

## **5.3 Preclinical safety data**

Apalutamide was negative for genotoxicity in a standard battery of *in vitro* and *in vivo* tests. Long-term animal studies have not been conducted to evaluate the carcinogenic potential of apalutamide.

Male fertility is likely to be impaired by treatment with apalutamide based on findings in repeat-dose toxicology studies which were consistent with the pharmacological activity of apalutamide. In repeat-dose toxicity studies in male rats and dogs, atrophy, aspermia/hypospermia, degeneration and/or hyperplasia or hypertrophy in the reproductive system were observed at doses corresponding to exposures approximately equal to the human exposure based on AUC.

In a fertility study in male rats, a decrease in sperm concentration and motility, copulation and fertility rates (upon pairing with untreated females) along with reduced weights of the secondary sex glands and epididymis were observed following 4 weeks of dosing at doses corresponding to exposures approximately equal to the human exposure based on AUC. Effects on male rats were reversible after 8 weeks from the last apalutamide administration.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

#### Tablet core

Colloidal anhydrous silica  
Croscarmellose sodium  
Hypromellose acetate succinate  
Magnesium stearate  
Microcrystalline cellulose  
Microcrystalline cellulose (silicified)

#### Film-coating

Iron oxide black (E172)  
Iron oxide yellow (E172)  
Macrogol  
Polyvinyl alcohol (partially hydrolysed)  
Talc  
Titanium dioxide (E171)

### **6.2 Incompatibilities**

Not applicable.

### **6.3 Shelf life**

2 years

#### **6.4 Special precautions for storage**

Store in the original package in order to protect from moisture.  
This medicinal product does not require any special temperature storage conditions.

#### **6.5 Nature and contents of container**

White opaque high-density polyethylene (HDPE) bottle with a polypropylene (PP) child-resistant closure. Each bottle contains 120 film-coated tablets and a total of 6 g of silica gel desiccant.

PVC-PCTFE foil blister with an aluminum push-through foil sealed inside a wallet pack.

- Each 28-day carton contains 112 film coated tablets in 4 cardboard wallet packs of 28 film-coated tablets each.
- Each 30-day carton contains 120 film coated tablets in 5 cardboard wallet packs of 24 film-coated tablets each.

Not all pack sizes may be marketed.

#### **6.6 Special precautions for disposal**

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

### **7. MARKETING AUTHORISATION HOLDER**

Janssen-Cilag International NV  
Turnhoutseweg 30  
B-2340 Beerse  
Belgium

### **8. MARKETING AUTHORISATION NUMBER(S)**

EU/1/18/1342/001  
EU/1/18/1342/002  
EU/1/18/1342/003

### **9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 14 January 2019

### **10. DATE OF REVISION OF THE TEXT**

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>.



**ANNEX II**

- A. MANUFACTURER(S) RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**
- C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**
- D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

**A. MANUFACTURER(S) RESPONSIBLE FOR BATCH RELEASE**

Janssen Cilag SpA  
Via C. Janssen  
Borgo San Michele  
Latina 04100, Italy

**B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**

Medicinal product subject to restricted medical prescription (see Annex I: Summary of Product Characteristics, section 4.2).

**C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**

• **Periodic safety update reports (PSURs)**

The requirements for submission of PSURs for this medicinal product are set out in the list of Union reference dates (EURD list) provided for under Article 107c(7) of Directive 2001/83/EC and any subsequent updates published on the European medicines web-portal.

The marketing authorisation holder (MAH) shall submit the first PSUR for this product within 6 months following authorisation.

**D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

• **Risk management plan (RMP)**

The marketing authorisation holder (MAH) shall perform the required pharmacovigilance activities and interventions detailed in the agreed RMP presented in Module 1.8.2 of the marketing authorisation and any agreed subsequent updates of the RMP.

An updated RMP should be submitted:

- At the request of the European Medicines Agency;
- Whenever the risk management system is modified, especially as the result of new information being received that may lead to a significant change to the benefit/risk profile or as the result of an important (pharmacovigilance or risk minimisation) milestone being reached.

• **Obligation to conduct post-authorisation measures**

The MAH shall complete, within the stated timeframe, the below measures:

<b>Description</b>	<b>Due date</b>
Postauthorisation efficacy study (PAES): In order to further evaluate the efficacy of Erleada, the MAH should submit the final clinical study report, including overall survival results, from study ARN-509-003 (SPARTAN) comparing the efficacy and safety of Apalutamide vs. placebo in subjects with high risk Non-Metastatic (M0) Castration-Resistant Prostate Cancer.	June 2023

**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**

## **A. LABELLING**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**CARTON 60 mg (BOTTLE)**

**1. NAME OF THE MEDICINAL PRODUCT**

Erleada 60 mg film-coated tablets  
apalutamide

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each film-coated tablet contains 60 mg apalutamide.

**3. LIST OF EXCIPIENTS**

**4. PHARMACEUTICAL FORM AND CONTENTS**

120 film-coated tablets

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Swallow the tablets whole.  
Read the package leaflet before use.  
Do not swallow or discard the desiccant.  
Oral use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

Store in original package in order to protect from moisture.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

Discard unused contents appropriately in accordance with local requirements.

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Janssen-Cilag International NV  
Turnhoutseweg 30  
B-2340 Beerse  
Belgium

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/18/1342/003

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

Erleada 60 mg

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

PC  
SN  
NN

**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGING**

**BOTTLE LABEL 60 mg**

**1. NAME OF THE MEDICINAL PRODUCT**

Erleada 60 mg film-coated tablets  
apalutamide

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each tablet contains 60 mg apalutamide.

**3. LIST OF EXCIPIENTS**

**4. PHARMACEUTICAL FORM AND CONTENTS**

120 tablets

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Swallow the tablets whole.  
Read the package leaflet before use.  
Oral use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

Store in original package in order to protect from moisture.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Janssen-Cilag International NV  
Turnhoutseweg 30  
B-2340 Beerse  
Belgium

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/18/1342/003

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

**17. UNIQUE IDENTIFIER – 2D BARCODE**

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**



**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**CARTON 60 mg (28 days)**

**1. NAME OF THE MEDICINAL PRODUCT**

Erleada 60 mg film-coated tablets  
apalutamide

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each film-coated tablet contains 60 mg apalutamide.

**3. LIST OF EXCIPIENTS**

**4. PHARMACEUTICAL FORM AND CONTENTS**

112 film-coated tablets

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Swallow the tablets whole.  
Read the package leaflet before use.  
Oral use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

Store in original package in order to protect from moisture.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

Discard unused contents appropriately in accordance with local requirements.

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Janssen-Cilag International NV  
Turnhoutseweg 30  
B-2340 Beerse  
Belgium

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/18/1342/001

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

Erleada 60 mg

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

PC  
SN  
NN

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**CARTON 60 mg (30 days)**

**1. NAME OF THE MEDICINAL PRODUCT**

Erleada 60 mg film-coated tablets  
apalutamide

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each film-coated tablet contains 60 mg apalutamide.

**3. LIST OF EXCIPIENTS**

**4. PHARMACEUTICAL FORM AND CONTENTS**

120 film-coated tablets

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Swallow the tablets whole.  
Read the package leaflet before use.  
Oral use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

Store in original package in order to protect from moisture.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

Discard unused contents appropriately in accordance with local requirements.

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Janssen-Cilag International NV  
Turnhoutseweg 30  
B-2340 Beerse  
Belgium

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/18/1342/002

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

Erleada 60 mg

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

PC  
SN  
NN

**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGING**

**OUTER WALLET 60 mg (28 days)**

**1. NAME OF THE MEDICINAL PRODUCT**

Erleada 60 mg film-coated tablets  
apalutamide

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each film-coated tablet contains 60 mg apalutamide.

**3. LIST OF EXCIPIENTS**

**4. PHARMACEUTICAL FORM AND CONTENTS**

28 film-coated tablets per wallet pack


**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Swallow the tablets whole.  
Read the package leaflet before use.  
Oral use.

(1) Press and hold



(2) Pull out



**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

Store in original package in order to protect from moisture.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

Discard unused contents appropriately in accordance with local requirements.

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Janssen-Cilag International NV  
Turnhoutseweg 30  
B-2340 Beerse  
Belgium

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/18/1342/001

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

Erleada 60 mg

**17. UNIQUE IDENTIFIER – 2D BARCODE**

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGING**

**OUTER WALLET 60 mg (30 days)**

**1. NAME OF THE MEDICINAL PRODUCT**

Erleada 60 mg film-coated tablets  
apalutamide

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each film-coated tablet contains 60 mg apalutamide.

**3. LIST OF EXCIPIENTS**

**4. PHARMACEUTICAL FORM AND CONTENTS**

24 film-coated tablets per wallet pack

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Swallow the tablets whole.  
Read the package leaflet before use.  
Oral use.

(1) Press and hold



(2) Pull out



**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**



**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

Store in original package in order to protect from moisture.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

Discard unused contents appropriately in accordance with local requirements.

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Janssen-Cilag International NV  
Turnhoutseweg 30  
B-2340 Beerse  
Belgium

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/18/1342/002

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY****15. INSTRUCTIONS ON USE****16. INFORMATION IN BRAILLE**

Erleada 60 mg

**17. UNIQUE IDENTIFIER – 2D BARCODE****18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

**MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS**

**INNER WALLET 60 mg (28 days)**

**1. NAME OF THE MEDICINAL PRODUCT**

Erleada 60 mg film-coated tablets  
apalutamide

**2. NAME OF THE MARKETING AUTHORISATION HOLDER**

Janssen-Cilag International NV

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. OTHER**

Fold over to close



Flip open



Monday  
Tuesday  
Wednesday  
Thursday  
Friday  
Saturday  
Sunday

**MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS**

**INNER WALLET 60 mg (30 days)**

**1. NAME OF THE MEDICINAL PRODUCT**

Erleada 60 mg film-coated tablets  
apalutamide

**2. NAME OF THE MARKETING AUTHORISATION HOLDER**

Janssen-Cilag International NV

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. OTHER**

Fold over to close



Flip open



Fill in your weekdays

Start date:

Day

**MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS**

**BLISTER 60 mg (12 count) (Blister sealed in inner wallet)**

**1. NAME OF THE MEDICINAL PRODUCT**

Erleada 60 mg film-coated tablets  
apalutamide

**2. NAME OF THE MARKETING AUTHORISATION HOLDER**

Janssen-Cilag International NV

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. OTHER**

**MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS**

**BLISTER 60 mg (16 count) (Blister sealed in inner wallet)**

**1. NAME OF THE MEDICINAL PRODUCT**

Erleada 60 mg film-coated tablets  
apalutamide

**2. NAME OF THE MARKETING AUTHORISATION HOLDER**

Janssen-Cilag International NV

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. OTHER**

**B. PACKAGE LEAFLET**

## Package leaflet: Information for the user

### Erleada 60 mg film-coated tablets apalutamide

▼ This medicine is subject to additional monitoring. This will allow quick identification of new safety information. You can help by reporting any side effects you may get. See the end of section 4 for how to report side effects.

#### **Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

#### **What is in this leaflet**

1. What Erleada is and what it is used for
2. What you need to know before you take Erleada
3. How to take Erleada
4. Possible side effects
5. How to store Erleada
6. Contents of the pack and other information

#### **1. What Erleada is and what it is used for**

Erleada is a cancer medicine that contains the active substance apalutamide.

It is used to treat adult men with prostate cancer that:

- has metastasised to other parts of the body and still responds to medical or surgical treatments that lower testosterone (also called hormone-sensitive prostate cancer).
- has not metastasised to other parts of the body and no longer responds to medical or surgical treatment that lowers testosterone (also called castration-resistant prostate cancer).

Erleada works by blocking the activity of hormones called androgens (such as testosterone). Androgens can cause the cancer to grow. By blocking the effect of androgens, apalutamide stops prostate cancer cells from growing and dividing.

#### **2. What you need to know before you take Erleada**

##### **Do not take Erleada if:**

- you are allergic to apalutamide or any of the other ingredients of this medicine (listed in section 6).
- you are a woman who is pregnant or may become pregnant (see the Pregnancy and contraception section below for more information).

Do not take this medicine if any of the above apply to you. If you are not sure, talk to your doctor or pharmacist before taking this medicine.

## **Warnings and precautions**

Talk to your doctor or pharmacist before taking this medicine if:

- you have ever had fits or seizures
- you are taking any medicines to prevent blood clots (e.g. warfarin, acenocoumarol)
- you have any heart or blood vessel conditions, including heart rhythm problems (arrhythmia).

Falls have been observed in patients taking Erleada. Take extra care to reduce your risk of a fall.

Broken bones have been observed in patients taking Erleada.

Blockage of the arteries in the heart that can lead to death has happened in some people during treatment with Erleada. Your healthcare provider will monitor you for signs and symptoms of heart problems during your treatment with Erleada. Call your healthcare provider or go to the nearest emergency room right away if you get chest pain or discomfort at rest or with activity or shortness of breath during your treatment with Erleada.

If you are taking any medicines, talk to your doctor or pharmacist to see if they are associated with an increased risk of seizure, bleeding or heart condition.

If any of the above apply to you (or you are not sure), talk to your doctor or pharmacist before taking Erleada.

## **Children and adolescents**

This medicine is not for use in children and adolescents under 18 years of age.

If a child or young person accidentally takes Erleada:

- go to the hospital straight away
- take this package leaflet with you to show to the emergency doctor.

## **Other medicines and Erleada**

Tell your doctor or pharmacist if you are taking, have recently taken, or might take any other medicines. This is because Erleada can affect the way some other medicines work. Also, some other medicines can affect the way Erleada works.

Tell your doctor if you are taking medicines that:

- lower high fat levels in the blood (e.g. gemfibrozil)
- treat bacterial infections (e.g. moxifloxacin, clarithromycin)
- treat fungal infections (e.g. itraconazole, ketoconazole)
- treat HIV infection (e.g. ritonavir, efavirenz, darunavir)
- treat anxiety (e.g. midazolam, diazepam)
- treat epilepsy (e.g. phenytoin, valproic acid)
- treat gastroesophageal reflux disease (conditions where there is too much acid in the stomach) (e.g. omeprazole)
- prevent blood clots (e.g. warfarin, clopidogrel, dabigatran etexilate)
- treat hayfever and allergies (e.g. fexofenadine)
- lower cholesterol levels (e.g. 'statins' such as rosuvastatin, simvastatin)
- treat heart conditions or lower blood pressure (e.g. digoxin, felodipine)
- treat heart rhythm problems (e.g. quinidine, disopyramide, amiodarone, sotalol, dofetilide, ibutilide)
- treat thyroid conditions (e.g. levothyroxine)
- treat gout (e.g. colchicine)
- lower blood glucose (e.g. repaglinide)
- treat cancer (e.g. lapatinib, methotrexate)
- treat opioid addiction or pain (e.g. methadone)
- treat serious mental illnesses (e.g. haloperidol)



You need to list the names of the medicines you take and show the list to your doctor or pharmacist when you start a new medicine. Mention to your doctor that you are taking Erleada if the doctor wants to start you on any new medicine. The dose of Erleada or any other medicines that you are taking may need to be changed.

### **Pregnancy and contraception information for men and women**

#### **Information for women**

- Erleada must not be taken by women who are pregnant, may become pregnant, or who are breast-feeding. Erleada may harm your unborn baby.

#### **Information for men – follow this advice during treatment and for 3 months after stopping**

- If you are having sex with a pregnant woman – use a condom to protect the unborn baby.
- If you are having sex with a woman who can become pregnant - use a condom and another highly effective method of contraception.

Use contraception during treatment and for 3 months after stopping. Talk to your doctor if you have any questions about contraception.

Erleada may reduce male fertility.

#### **Driving and using machines**

This medicine is not likely to affect you being able to drive and use any tools or machines. The side effects for Erleada include seizures. If you are at higher risk of seizures (see Section 2 Warnings and precautions), talk to your doctor.

#### **Erleada contains sodium**

This medicine contains less than 1 mmol sodium (23 mg) per 240 mg dose (4 tablets), that is to say essentially 'sodium-free'.

### **3. How to take Erleada**

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

#### **How much to take**

The recommended dose is 240 mg (four 60 mg tablets) once a day.

#### **Taking Erleada**

- Take this medicine by mouth.
- You can take Erleada with food or between meals.
- Swallow the tablets whole.

Your doctor may also prescribe other medicines while you are taking Erleada.

#### **If you take more Erleada than you should**

If you take more than you should, stop taking Erleada and contact your doctor. You may have an increased risk of side effects.

#### **If you forget to take Erleada**

If you forget to take Erleada, take your usual dose as soon as you remember.

- If you forget to take Erleada for the whole day - take your usual dose the following day.
- If you forget to take Erleada for more than one day - talk to your doctor straight away.

Do not take a double dose to make up for a forgotten dose.

**If you stop taking Erleada**

Do not stop taking Erleada without checking with your doctor first.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

**4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

**Stop using Erleada and seek medical attention immediately if you notice any of the following symptoms:**

- reddish non-elevated, target-like or circular patches on the trunk, often with central blisters, skin peeling, ulcers of mouth, throat, nose, genitals and eyes. These serious skin rashes can be preceded by fever and flu-like symptoms (toxic epidermal necrolysis).

**Serious side effects**

Tell your doctor straight away if you notice any of the following serious side effects – your doctor may stop treatment:

- fit or seizure – this is uncommon (may affect up to 1 in 100 people). Your healthcare provider will stop Erleada if you have a seizure during treatment.
- falls or fractures (broken bones) – these are very common (may affect more than 1 in 10 people). Your healthcare provider may monitor you more closely if you are at risk for fractures.
- heart disease – this is common (may affect up to 1 in 10 people). Your healthcare provider will monitor you for signs and symptoms of heart problems during your treatment. Call your healthcare provider or go to the nearest emergency room right away if you get chest pain or discomfort at rest or with activity or shortness of breath during your treatment with Erleada.

Tell your healthcare provider right away if you notice any of the serious side effects above.

**Other side effects include**

**Very common** (may affect more than 1 in 10 people):

- feeling very tired
- joint pain
- skin rash
- decreased appetite
- high blood pressure
- hot flush
- diarrhoea
- broken bones
- falls
- weight loss.

**Common** (may affect up to 1 in 10 people):

- muscle spasms
- itching
- change in sense of taste
- blood test showing high level of cholesterol in the blood
- blood test showing high level of a type of fat called “triglycerides” in the blood
- heart disease
- under-active thyroid which can make you feel more tired and have difficulty getting started in the morning, and blood tests may also show an under-active thyroid.

**Not known** (frequency cannot be estimated from the available data):

- abnormal heart tracing on an ECG (electrocardiogram)

- life-threatening rash with blisters and peeling over much of the body (toxic epidermal necrolysis).

### **Reporting of side effects**

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system](#) listed in [Appendix V](#). By reporting side effects, you can help provide more information on the safety of this medicine.

## **5. How to store Erleada**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the container (blister foils, inner wallet, outer wallet, bottle, and carton) after EXP. The expiry date refers to the last day of that month.

Store in the original package in order to protect from moisture. This medicine does not require any special temperature storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## **6. Contents of the pack and other information**

### **What Erleada contains**

- The active substance is apalutamide. Each film-coated tablet contains 60 mg of apalutamide.
- The other ingredients of the tablet core are colloidal anhydrous silica, croscarmellose sodium, hypromellose acetate succinate, magnesium stearate, microcrystalline cellulose, and silicified microcrystalline cellulose. The film-coating contains iron oxide black (E172), iron oxide yellow (E172), macrogol, polyvinyl alcohol (partially hydrolysed), talc, and titanium dioxide (E171).

### **What Erleada looks like and contents of the pack**

Erleada film-coated tablets are slightly yellowish to greyish green, oblong-shaped, film-coated tablets (16.7 mm long x 8.7 mm wide), with “AR 60” written on one side.

The tablets may be supplied either in a bottle or in a wallet pack. Not all pack sizes may be marketed.

### **Bottle**

The tablets are supplied in a plastic bottle with a child-resistant closure. Each bottle contains 120 tablets and a total of 6 g of desiccant. Each carton contains one bottle. Store in the original package. Do not swallow or discard desiccant.

### **28-day carton**

Each 28-day carton contains 112 film-coated tablets in 4 cardboard wallet packs of 28 film-coated tablets each.

### **30-day carton**

Each 30-day carton contains 120 film-coated tablets in 5 cardboard wallet packs of 24 film-coated tablets each.

### **Marketing Authorisation Holder**

Janssen-Cilag International NV  
Turnhoutseweg 30

B-2340 Beerse  
Belgium

**Manufacturer**

Janssen Cilag SpA  
Via C. Janssen  
Borgo San Michele  
Latina 04100, Italy

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

**België/Belgique/Belgien**

Janssen-Cilag NV  
Tel/Tél: +32 14 64 94 11  
janssen@jacbe.jnj.com

**България**

„Джонсън & Джонсън България” ЕООД  
Тел.: +359 2 489 94 00  
jjsafety@its.jnj.com

**Česká republika**

Janssen-Cilag s.r.o.  
Tel.: +420 227 012 227

**Danmark**

Janssen-Cilag A/S  
Tlf: +45 4594 8282  
jacdk@its.jnj.com

**Deutschland**

Janssen-Cilag GmbH  
Tel: +49 2137 955 955  
jancil@its.jnj.com

**Eesti**

UAB "JOHNSON & JOHNSON" Eesti filiaal  
Tel: +372 617 7410  
ee@its.jnj.com

**Ελλάδα**

Janssen-Cilag Φαρμακευτική Α.Ε.Β.Ε.  
Τηλ: +30 210 80 90 000

**España**

Janssen-Cilag, S.A.  
Tel: +34 91 722 81 00  
contacto@its.jnj.com

**France**

Janssen-Cilag  
Tél: 0 800 25 50 75 / +33 1 55 00 40 03  
medisource@its.jnj.com

**Lietuva**

UAB "JOHNSON & JOHNSON"  
Tel: +370 5 278 68 88  
lt@its.jnj.com

**Luxembourg/Luxemburg**

Janssen-Cilag NV  
Tél/Tel: +32 14 64 94 11  
janssen@jacbe.jnj.com

**Magyarország**

Janssen-Cilag Kft.  
Tel.: +36 1 884 2858  
janssenhu@its.jnj.com

**Malta**

AM MANGION LTD.  
Tel: +356 2397 6000

**Nederland**

Janssen-Cilag B.V.  
Tel: +31 76 711 1111  
janssen@jacnl.jnj.com

**Norge**

Janssen-Cilag AS  
Tlf: +47 24 12 65 00  
jacno@its.jnj.com

**Österreich**

Janssen-Cilag Pharma GmbH  
Tel: +43 1 610 300

**Polska**

Janssen-Cilag Polska Sp. z o.o.  
Tel.: +48 22 237 60 00

**Portugal**

Janssen-Cilag Farmacêutica, Lda.  
Tel: +351 214 368 600

**Hrvatska**

Johnson & Johnson S.E. d.o.o.  
Tel: +385 1 6610 700  
jjsafety@JNJCR.JNJ.com

**Ireland**

Janssen Sciences Ireland UC  
Tel: +353 1 800 709 122

**Ísland**

Janssen-Cilag AB  
c/o Vistor hf.  
Sími: +354 535 7000  
janssen@vistor.is

**Italia**

Janssen-Cilag SpA  
Tel: 800.688.777 / +39 02 2510 1  
janssenita@its.jnj.com

**Κύπρος**

Βαρνάβας Χατζηπαναγής Λτδ  
Τηλ: +357 22 207 700

**Latvija**

UAB "JOHNSON & JOHNSON" filiāle Latvijā  
Tel: +371 678 93561  
lv@its.jnj.com

**România**

Johnson & Johnson România SRL  
Tel: +40 21 207 18 00

**Slovenija**

Johnson & Johnson d.o.o.  
Tel: +386 1 401 18 00  
Janssen\_safety\_slo@its.jnj.com

**Slovenská republika**

Johnson & Johnson, s.r.o.  
Tel: +421 232 408 400

**Suomi/Finland**

Janssen-Cilag Oy  
Puh/Tel: +358 207 531 300  
jacfi@its.jnj.com

**Sverige**

Janssen-Cilag AB  
Tfn: +46 8 626 50 00  
jacse@its.jnj.com

**United Kingdom**

Janssen-Cilag Ltd.  
Tel: +44 1 494 567 444

**This leaflet was last revised in MM/YYYY.**

**Other sources of information**

Detailed information on this medicine is available on the European Medicines Agency web site:  
<http://www.ema.europa.eu>